



The selective p38 inhibitor SB-239063 protects primary neurons from mild to moderate excitotoxic injury

Jeffrey J. Legos^{a,*}, BethAnn McLaughlin^b, Stephen D. Skaper^c, Paul J.L.M. Strijbos^c, Andrew A. Parsons^c, Elias Aizenman^b, Greta A. Herin^b, Frank C. Barone^a, Joseph A. Erhardt^d

^a*In Vivo Pharmacology, High Throughput Biology, Discovery Research, GlaxoSmithKline, PO Box 1539, Mail Code UW 2523, 709 Swedeland Road, King of Prussia, PA 19406, USA*

^b*Department of Neurobiology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA*

^c*Neurology Centre of Excellence for Drug Discovery, GlaxoSmithKline, Harlow, UK*

^d*Cardiovascular and Urogenital Center of Excellence for Drug Discovery, GlaxoSmithKline, King of Prussia, PA, USA*

Received 5 April 2002; received in revised form 23 May 2002; accepted 24 May 2002

Abstract

Inhibition of the p38 mitogen-activated protein kinase (MAP Kinase) pathway reduces acute ischemic injury in vivo, suggesting a direct role for this signaling pathway in a number of neurodegenerative processes. The present study was designed to evaluate further the role of p38 MAP Kinase in acute excitotoxic neuronal injury using the selective p38 inhibitor SB-239063 (*trans*-1-(4-hydroxycyclohexyl)-4-(fluorophenyl)-5-(2-methoxy-pyrimidin-4-yl) imidazole). Unlike the widely used p38 inhibitor, SB-203580 (4-(4-Fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)1*H*-imidazole), this second generation p38 inhibitor more selectively inhibits p38 MAP Kinase without affecting the activity of other MAP Kinase signaling pathways and provides a more accurate means to selectively assess the role of p38 in excitotoxicity that has not been previously possible. SB-239063 provided substantial protection against cell death induced by either oxygen glucose deprivation (OGD) or magnesium deprivation in cultured neurons. The ability of this compound to block excitotoxicity was not due to direct inhibition of *N*-methyl-D-aspartate (NMDA) receptor-mediated currents as SB-239063 did not alter NMDA electrophysiological responses. SB-239063 did not protect against a severe excitotoxic insult induced by 60-min exposure to NMDA. However, when tested against a less severe, brief (5 min) NMDA exposure, p38 inhibition provided substantial protection. These data demonstrate that inhibition of p38 MAP Kinase can confer neuroprotection in vitro against mild but not severe excitotoxic exposure, and suggests that other additional pathways/mechanism(s) may be involved in severe excitotoxic cell death. © 2002 Published by Elsevier Science B.V.

Keywords: p38 MAP Kinase; Acute neuronal injury; Cell culture; Neuroprotection; Ischemic cell death; Excitotoxicity

1. Introduction

The mitogen-activated protein kinase family (MAP Kinase) regulate cell stress and survival-related signals by phosphorylating intracellular enzymes and transcription factors (Cobb, 1999). Specific MAP Kinases are involved in cell survival, apoptosis, inflammatory cytokine production, and possibly secondary ischemia (Barone and Parsons, 2000; English et al., 1999; McLaughlin et al., 2001). However, the exact role(s) of MAP Kinases in mediating excitotoxic central nervous system (CNS) injury have not been fully elucidated. While a number of reports have suggested a critical role for p38 activation following *N*-methyl-D-aspar-

tate (NMDA) receptor overstimulation (Kawasaki et al., 1997; Stanciu et al., 2000), these studies have been restricted to the use of less selective inhibitors which can effect other MAP Kinase signaling pathways. Recently, the selective p38 MAP Kinase inhibitor, SB-239063 (*trans*-1-(4-hydroxycyclohexyl)-4-(fluorophenyl)-5-(2-methoxy-pyrimidin-4-yl) imidazole), has been shown to provide significant neuroprotection in rodent focal stroke (Barone et al., 2001a,b; Legos et al., 2001). The aim of the present study was to further evaluate the direct neuroprotective effects of p38 MAP Kinase inhibition in various in vitro neuronal cell models of excitotoxic injury. This clarification of the direct contribution of p38 to excitotoxicity should improve our understanding of the mechanism(s) involved in ischemia-induced brain injury, and hopefully lead to more effective therapeutic strategies in ischemic and traumatic brain injury (Barone and Parsons, 2000; Lee and Young, 1996).

* Corresponding author. Tel.: +1-610-270-6547; fax: +1-610-270-4114.
E-mail address: Jeffrey_J_Legos@gsk.com (J.J. Legos).

59 2. Materials and methods

60 2.1. Preparation of neuron-enriched forebrain cultures

62 Neuron-enriched forebrain cultures were prepared from
63 embryonic day 17 (E17) rat fetuses as previously described
64 (McLaughlin et al., 1998). Dissociated cells were plated on
65 poly-L-ornithine-treated tissue culture plates in a growth
66 medium comprised of 80% Dulbecco's modified eagle's me-
67 dium (DMEM) (high glucose with L-glutamine but without
68 sodium pyruvate), 10% Ham's F-12 nutrients, 10% bovine
69 calf serum (heat-inactivated) with antimycotic/antibiotic
70 mixture (with amphotericin B and streptomycin sulfate).
71 Cultures were maintained in an incubator at 37 °C, 95%
72 air/5% CO₂. Glial cell proliferation was inhibited after 48 h in
73 culture with 1–2 μM cytosine arabinoside. After 3 days in
74 vitro, medium was replaced with a serum-free solution com-
75 prised of Neurobasal medium (without L-glutamine), B27
76 supplement, and antimycotic/antibiotic mixture. At 2 weeks
77 in vitro, these cultures were >95% neuronal as assessed by
78 Hoechst and glial fibrillary acidic protein staining.

79 2.2. Oxygen glucose deprivation injury

81 Oxygen glucose deprivation (OGD) was performed on 2-
82 week-old cultures. Sixty minutes prior to ischemia, cells
83 were rinsed in Minimal Eagle's Medium with Earle's salts
84 solution, previously saturated with 10% H₂/85% N₂/5%
85 CO₂. Cultures were kept in an anaerobic chamber for various
86 times (45, 60, 120 min) at 37 °C and anoxia–ischemia was
87 terminated by replacement of the Earle's balanced salt
88 solution with oxygenated growth medium containing vehicle
89 or SB-239063 (20 μM). Cell viability was assessed 20–24 h
90 after oxygen glucose deprivation by measuring the extent of
91 lactate dehydrogenase (LDH) release into the medium as
92 previously described (Hartnett et al., 1997). Media samples
93 (40 μl) were analysed spectrophotometrically (490:630 nm)
94 according to the manufacturer's instructions.

95 2.3. Preparation of mixed cortical cultures for electro- 96 physiology and NMDA exposure

98 Electrophysiology and toxicity experiments assessing the
99 effects of SB-239063 on NMDA receptors were performed in
100 mixed forebrain cultures of neurons and glia. Cerebral
101 cortices were obtained from embryonic day 16 (E16) Sprague
102 Dawley rat fetuses and dissociated as previously described
103 (Hartnett et al., 1997). Briefly, cells were plated onto poly-L-
104 lysine-coated glass coverslips at a density of 225,000 cells/ml
105 of growth medium (v/v mixture of 80% DMEM, 10% Ham's
106 F12, 10% calf serum, 25 mM HEPES, 24 U/ml penicillin, 24
107 μg/ml streptomycin, and 2 mM L-glutamine) and maintained
108 at 37 °C in 95% air/5% CO₂. Cytosine arabinoside (2 μM)
109 was added once at 15 days in vitro after which growth
110 medium lacking F-12 and containing low serum (2%) was
111 added. Medium was partially replaced with fresh growth

medium three times per week. At 3–5 weeks in vitro, these
cultures contain ~10–20% neurons (Rosenberg, 1991;
Rosenberg and Aizenman, 1989).

112 2.4. Electrophysiological recordings

117 Electrophysiology experiments were performed at room
118 temperature (25 °C) using the whole-cell patch clamp con-
119 figuration. Coverslips were bathed in external solution con-
120 taining (concentrations expressed in mM): 150 NaCl, 1.0
121 CaCl₂, 2.8 KCl, 10 HEPES, 10 glycine, 25 tetrodotoxin
122 (Calbiochem) and pH was adjusted to 7.2 with NaOH.
123 Electrodes were pulled on a Sutter P-87 electrode puller
124 (Sutter Instruments, Novato, CA) to a resistance of 1.5–3
125 MΩ when filled with internal solution containing (in mM):
126 140 CsF, 10 EGTA/CsOH, 1 CaCl₂, and 10 HEPES (pH
127 adjusted to 7.2 with CsOH). Signals were amplified using an
128 Axopatch 200B integrating patch clamp amplifier (Axon
129 Instruments, Foster City, CA), filtered at 1 kHz, and digitised
130 at 2 kHz with a DigiData 1200 (Axon Instruments) computer
131 interface. Drugs were applied via a perfusion system with a
132 stepper motor for fast solution changes (Warner Instruments,
133 Hamden, CT). NMDA and SB-239063 were dissolved in
134 external solution for recording. SB-239063 was diluted from
135 a 1000× stock. Data were collected and analysed using
136 commercially available software (pCLAMP 6.11, Axon
137 Instruments).

138 2.5. Preparation of hippocampal cell cultures

140 Neuron-enriched hippocampal cultures were prepared
141 from embryonic Sprague–Dawley rat fetuses (gestational
142 age 17.5 days; Charles River) as described previously
143 (Skaper et al., 2001). Hippocampi were incubated with
144 0.08% (w/v) trypsin, and dissociated in Neurobasal medium
145 containing 10% heat-inactivated fetal calf serum (Skaper et
146 al., 1990). Cells were pelleted by centrifugation (200×g, 5
147 min) and resuspended in Neurobasal medium containing
148 B27 supplement, 25 μM glutamate, 1 mM sodium pyruvate,
149 2 mM L-glutamine, 50 U/ml penicillin, and 50 μg/ml
150 streptomycin. The cell suspension was plated onto dishes
151 previously coated with poly-D-lysine (10 μg/ml) and 10%
152 heat-inactivated fetal calf serum, at a density of 4.5×10⁴
153 cells/cm². Cultures were maintained at 37 °C in a humidified
154 atmosphere of 5% CO₂/95% air. After 5 days, one-half the
155 medium was replaced with an equal volume of maintenance
156 medium (plating medium but containing B27 supplements
157 without antioxidants, and lacking glutamate). Additional
158 medium exchanges (0.5 volume) were performed every 3–
159 4 days thereafter.

160 2.6. Magnesium withdrawal treatment to generate excito- 161 toxicity

163 Toxicity experiments were performed on cells between 14
164 and 16 days in vitro. Cultures were washed once with

165 $Mg^{2+}Cl_2$ -free Locke's solution (pH 7.0) containing 0.1 μM
 166 glycine and 30 μM histamine (Skaper et al., 2001). Control
 167 cultures were exposed to Locke's solution containing 1 mM
 168 $Mg^{2+}Cl_2$. Drug treatments were carried out for 15 min (22
 169 $^{\circ}C$) in a final volume of 0.5 ml. Thereafter, cells were
 170 washed with complete Locke's solution and returned to their
 171 original culture medium for 24 h. Cell survival was quanti-
 172 fied 24 h after the insult by a colorimetric reaction with 3-
 173 (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
 174 (MTT) (Skaper et al., 1990). Absolute MTT values obtained
 175 were normalized and expressed as a percentage of sham-
 176 treated sister cultures (defined as 100%).

177

178 2.7. NMDA exposure in mixed cultures

179 Excitotoxicity generated by NMDA exposure was per-
 180 formed in 4-week-old mixed cultures of neurons and glia,
 181 prepared as described for electrophysiological recordings.
 182 Cell cultures were pretreated for 1 h with SB-239063 (20
 183 μM), followed by a 5-min or 1-h exposure to NMDA in the
 184 presence of 10 μM glycine. SB-239063 was present through-
 185 out the treatment and over the subsequent 20–24 h until LDH
 186 readings were performed.

187 3. Results

188

189 3.1. SB-239063 protects against oxygen glucose deprivation

190 Based on the dose–response curve obtained in prelimi-
 191 nary experiments (data not shown), the effects of SB-239063
 192 (20 μM) on oxygen glucose deprivation toxicity were eval-
 193 uated at various time points in neuron-enriched forebrain
 194 cultures. Cells were exposed for 45, 60, or 120 min to oxygen
 195 glucose deprivation in the presence or absence of the p38
 196 inhibitor (Fig. 1). Forty-five minutes of oxygen glucose
 197 deprivation did not significantly increase the amount of
 198 LDH released (0.016 ± 0.0003 optical density units, $n=3$)
 199 compared to control (0.010 ± 0.001 optical density units,
 200 $n=3$). However, following 60 min of oxygen glucose depri-
 201 vation, appreciable cell death was present and SB-239063
 202 significantly ($P < 0.05$) decreased LDH release (0.023 ± 0.004
 203 optical density units, $n=3$) compared to vehicle treatment
 204 (0.041 ± 0.005 optical density units, $n=3$). Although the
 205 extent of cell death after 60 min of oxygen glucose depriva-
 206 tion was similar to that seen after 120 min of oxygen glucose
 207 deprivation (0.038 ± 0.006 optical density units, $n=3$), SB-
 208 239063 did not protect against the latter treatment ($0.037 \pm$
 209 0.007 optical density units, $n=3$), suggesting that this dura-
 210 tion of oxygen glucose deprivation was too severe to permit
 211 pharmacological intervention.

212

213 3.2. SB-239063 does not block NMDA-mediated currents

214 In order to insure that any neuroprotective action of SB-
 215 239063 was not due to a direct effect on NMDA receptor-




Fig. 1. (A) SB-239063 blocks excitotoxicity induced by oxygen glucose deprivation in neuron-enriched forebrain cultures. Neuronal cell cultures were exposed for various times to oxygen glucose deprivation. Addition of SB-239063 (20 μM) significantly decreased neuronal cell death following 60 min of oxygen glucose deprivation. Data are means \pm S.E.M. and were analysed by two-tailed paired *t*-test (significance at * $P < 0.05$). Representative photomicrographs were taken of cell cultures 24 h after 60-min oxygen glucose deprivation treated with either vehicle (B) or 20 μM SB-239063 (C). Cells which had lost their phase-bright appearance become shrunken (pyknotic) and were dying following treatment are indicated by black arrowheads. Scale bar is 75 μm .

mediated channel activity, whole cell electrophysiological
 recordings were performed. In these experiments, 30 μM
 NMDA was applied to mixed cultures of neurons and glia in
 the presence and absence of SB-239063 (Fig. 2). The p38
 inhibitor did not alter NMDA elicited currents suggesting
 that any neuroprotective action of the MAP Kinase inhibitor
 was not a direct effect on NMDA receptors.

216
 217
 218
 219
 220
 221
 222




Fig. 2. SB-239063 does not alter NMDA-elicited electrophysiological responses. A representative trace of whole-cell responses to NMDA in the presence and absence of SB-239063 (20 μ M). Application of SB-239063 was made for 3 s concomitantly during exposure to 30 μ M NMDA. The p38 MAP Kinase inhibitor did not alter responses to agonist in any of the cells from which recordings were made ($n=4$).

223

224 3.3. Excitotoxicity generated by magnesium withdrawal can
225 be attenuated with SB-239063

226 We used neuron-enriched hippocampal cell cultures to
227 assess the effects of p38 inhibition on moderate excitotoxi-
228 city generated by removal of magnesium block from the
229 NMDA receptor. Control experiments showed that the loss
230 of viable neurons, as quantified by MTT assay, was propor-
231 tional to the number of degenerating neurons, as estimated
232 by trypan blue staining. By 24 h, approximately 50% of
233 neurons treated with vehicle were no longer viable as
234 assessed by MTT ($52.0 \pm 5.6\%$). Treatment with SB-
235 239063 (1–30 μ M) provided dose-related reduction in




Fig. 3. SB-239063 reduces excitotoxic hippocampal neuronal cell death. In dissociated hippocampal cells, SB-239063 (10–30 μ M) decreased neuronal cell death by 25–43% following Mg^{2+} withdrawal. Absolute MTT values obtained were normalized and expressed as a percentage of sham-treated sister cultures (defined as 100% survival). Percent cell death was calculated from these data. Values are means \pm S.D. ($n=3$) and were analysed by ANOVA followed by Dunnett's post-hoc test (significance at $*P<0.05$).

neuronal cell death (Fig. 3). SB-239063 was maximally
neuroprotective at 10–30 μ M with neuronal survival sig-
nificantly ($P<0.05$) increasing by 25–43% ($35.8 \pm 2.4\%$ and
 $27.7 \pm 2.4\%$).

236
237
238
239
240
241
242

3.4. SB-239063 reduces cell death associated with mild, but
not severe, NMDA exposure

As our earlier OGD experiments suggested that p38
inhibition may be more efficacious with mild excitotoxic
insults, we next assessed the relative contribution of p38 to
direct exposure to NMDA. In these studies, we assessed the
efficacy of SB-239063 at attenuating mild (5 min) and severe
(1 h) excitotoxicity in mixed cultures of neurons and glia.
Cultures that were exposed for 1 h to either 30, 100 or 300
 μ M NMDA were not significantly protected by 20 μ M SB-
239063. However, when cells were exposed to a more mild
insult (5 min), SB-239063 (20 μ M) was capable of attenuat-
ing NMDA induced injury. Statistically significant neuro-


243
244
245
246
247
248
249
250
251
252
253

Fig. 4. SB-239063 is protective against mild, but not severe, NMDA exposure in mixed cortical cell cultures. (A) Four-week-old mixed cortical cell cultures were treated with vehicle or SB 239063 (20 μ M) 1 h prior to exposure to NMDA (30, 100, and 300 μ M). NMDA was removed after 1 h and cell death was measured 24 h later using an LDH assay. Data are means \pm S.D. ($n=6$) and analysed by ANOVA. (B) Paired 4-week-old mixed cultures were exposed to various concentrations of NMDA (0.01, 0.03, 0.1, 0.3, 1, and 3 mM) in the presence or absence of SB-239063 as above except NMDA was removed after 5-min incubation. Data are means \pm S.E.M. ($n=5-7$) and were analysed by paired Student's t -test (significance at $*P<0.05$).

254 protection was observed at 100 μ M, 300 μ M, 1 mM and 3
255 mM NMDA with SB-239063 providing between 20% and
256 28% protection at these concentrations (Fig. 4).

257 4. Discussion

258 Several groups have reported that both glutamate and
259 hypoxia activate p38 in neuronal cell cultures (Clerk et al.,
260 1998; Kawasaki et al., 1997). However, in assessing a role for
261 this MAP Kinase in the observed toxicity, these studies have
262 relied on less specific MAP Kinase inhibitors such as SB-
263 203580 (4-(4-Fluorophenyl)-2-(4-methylsulfinylphenyl)-5-
264 (4-pyridyl)1*H*-imidazole). The use of SB-239063 in the
265 current study provides a number of advantages over SB-
266 203580. We demonstrate in this work that SB-239063 does
267 not directly alter NMDA induced currents, and we have
268 previously shown that SB-239063 does not effect other
269 MAP Kinases. SB-203580, however, has been shown to
270 inhibit c-Jun N-terminal kinase (JNK) activity with an IC_{50}
271 of 5 μ M and cRaf with an IC_{50} of 0.4 μ M (Barone et al.,
272 2001a). Thus, SB-239063 provides a more precise means to
273 assess the role of p38 in excitotoxicity. Indeed, this is the first
274 study to unequivocally demonstrate a role for p38 in excito-
275 toxicity. In this work, we demonstrates that activation of p38
276 MAP Kinase plays a critical role in neuronal cell death
277 induced by mild oxygen glucose deprivation, magnesium
278 withdrawal and glutamate receptor agonist exposure. These
279 models share a requirement for NMDA receptor activation,
280 (Kaku et al., 1991; Skaper et al., 2001; Speliotos et al., 1994)
281 and are therefore considered excitotoxic in nature.

282 In previous studies, we and others have shown that several
283 intracellular signals commonly associated with glutamate
284 receptor stimulation and excitotoxicity may contribute to
285 p38 activation. For instance, Koh et al. (1996) demonstrated
286 that zinc is released during transient ischemia and ‘free’ zinc
287 accumulates in dying neurons. We have recently shown that
288 oxidant-induced zinc dysregulation is the most proximal
289 events in an apoptotic cascade in which p38 activation leads
290 to potassium efflux, and subsequently, energetic dysfunction
291 and caspase activation (McLaughlin et al., 2001). Blockade
292 of p38 with SB-239063 provided substantial neuroprotection
293 against this oxidative and ionic dysfunction. Given the
294 importance of oxidative stress and zinc dysregulation in
295 excitotoxic insults, our current observations are perhaps not
296 surprising as they closely parallel our previous observations.
297 p38 has been shown to be a critical mediator of the
298 inflammatory response in CNS (Irving et al., 2000). Earlier
299 work from our labs has shown SB-239063 can attenuate
300 early neuronal injury (within 2 h) in an animal model of
301 cerebral ischemia induced by electrocoagulation of the mid-
302 dle cerebral artery (Legos et al., 2001). This protection was
303 maintained for at least 7 days suggesting a direct, long-
304 lasting protective effect against the subsequent apoptotic cell
305 death which is typical of this model. As cytokine production
306 and neutrophil infiltration might be delayed up to 12 h

(Legos et al., 2000), these initial studies demonstrating early
neuroprotection by SB-239063 suggests that p38 MAP
Kinase inhibitors can be beneficial via additional or alter-
native pathways which do not involve inflammation per se,
such as blockade of excitotoxic signaling.

Taken together, these data demonstrate that in addition to
its role in mediating the inflammatory response, p38 MAP
Kinase may also contribute to early neuronal injury through
excitotoxic pathways. It is, however, important to note that in
excitotoxic models where significant protection was afforded
by blocking p38, a substantial amount of cell death
remained. This argues that even mild excitotoxicity clearly
has additional p38-independent components which contrib-
ute to cell death. These experiments, in conjunction with
others (Barone et al., 2001a,b; Irving et al., 2000; Walton et
al., 1998), suggest that activation of p38 and other MAP
Kinases and their functional importance may be specific to
various cell types and/or duration and intensity of the
stimulus. Low level excitotoxic insults may activate apop-
totic signaling pathways in a manner similar to what we have
previously shown in a subpopulation of cells which are
responsive to p38 inhibition. However, more severe insults
may not require apoptotic signaling cascades to cause death,
are impervious to p38 blockade, and involve other signal
transduction cascades. Taken together, these studies suggest
that in instances of mild to moderate glutamatergic over-
activation, p38 MAP Kinase inhibition may provide a means
to prevent or minimize neuronal cell death. These results
may be applicable to neurological disorders that have been
associated with excitotoxic injury such as cardiac arrest,
stroke and epilepsy.

Acknowledgements

The authors would like to thank Karen Hartnett and Shen
Du for their technical assistance in performing some of these
experiments.

References

- Barone, F.C., Parsons, A.A., 2000. Therapeutic potential of anti-inflamma-
tory drugs in focal stroke. *Exp. Opin. Invest. Drugs* 9, 2281–2306.
- Barone, F.C., Irving, E.A., Ray, A.M., Lee, J.C., Kassis, S., Kumar, S.,
Badger, A.M., Legos, J.J., Erhardt, J.A., Ohlstein, E.H., Hunter, A.J.,
Harrison, D.C., Philpott, K., Smith, B.R., Adams, J.L., Parsons, A.A.,
2001a. Inhibition of p38 mitogen-activated protein kinase provides neu-
roprotection in cerebral focal ischemia. *Med. Res. Rev.* 21, 129–145.
- Barone, F.C., Irving, E.A., Ray, A.M., Lee, J.C., Kassis, S., Kumar, S.,
Badger, A.M., White, R.F., McVey, M.J., Legos, J.J., Erhardt, J.A., Nel-
son, A.H., Ohlstein, E.H., Hunter, A.J., Ward, K., Smith, B.R., Adams,
J.L., Parsons, A.A., 2001b. SB 239063, a second-generation p38 mito-
gen-activated protein kinase inhibitor, reduces brain injury and neuro-
logical deficits in cerebral focal ischemia. *J. Pharmacol. Exp. Ther.* 296,
312–321.
- Clerk, A., Fuller, S.J., Michael, A., Sugden, P.H., 1998. Stimulation of
“stress-regulated” mitogen-activated protein kinases (stress-activated
protein kinases/c-Jun N-terminal kinases and p38-mitogen-activated pro-

- tein kinases) in perfused rat hearts by oxidative and other stresses. *J. Biol. Chem.* 273, 7228–7234.
- Cobb, M.H., 1999. MAP kinase pathways. *Prog. Biophys. Mol. Biol.* 71, 479–500.
- English, J., Pearson, G., Wilsbacher, J., Swantek, J., Karandikar, M., Xu, S.C., Cobb, M.H., 1999. New insights into the control of MAP kinase pathways. *Exp. Cell Res.* 253, 255–270.
- Hartnett, K.A., Stout, A.K., Rajdev, S., Rosenberg, P.A., Reynolds, I.J., Aizenman, E., 1997. NMDA receptor-mediated neurotoxicity—a paradoxical requirement for extracellular Mg^{2+} in Na^+/Ca^{2+} -free solutions in rat cortical neurons in vitro. *J. Neurochem.* 68, 1836–1845.
- Irving, E.A., Barone, F.C., Reith, A.D., Hadingham, S.J., Parsons, A.A., 2000. Differential activation of MAPK/ERK and p38/SAPK in neurons and glia following focal cerebral ischaemia in the rat. *Mol. Brain Res.* 77, 65–75.
- Kaku, D.A., Goldberg, M.P., Choi, D.W., 1991. Antagonism of non-NMDA receptors augments the neuroprotective effect of NMDA receptor blockade in cortical cultures subjected to prolonged deprivation of oxygen and glucose. *Brain Res.* 554, 344–347.
- Kawasaki, H., Morooka, T., Shimohama, S., Kimura, J., Hirano, T., Gotoh, Y., Nishida, E., 1997. Activation and involvement of p38 mitogen-activated protein kinase in glutamate-induced apoptosis in rat cerebellar granule cells. *J. Biol. Chem.* 272, 18518–18521.
- Lee, J.C., Young, P.R., 1996. Role of CSB/p38/RK stress response kinase in LPS and cytokine signaling mechanisms. *J. Leukoc. Biol.* 59, 152–157.
- Legos, J.J., Whitmore, R.G., Erhardt, J.A., Parsons, A.A., Tuma, R.F., Barone, F.C., 2000. Quantitative changes in interleukin proteins following focal stroke in the rat. *Neurosci. Lett.* 282, 189–192.
- Legos, J.J., Erhardt, J.A., White, R.F., Lenhard, S.C., Chandra, S., Parsons, A.A., Tuma, R.F., Barone, F.C., 2001. SB 239063, a novel p38 inhibitor, attenuates early neuronal injury following ischemia. *Brain Res.* 892, 70–77.
- McLaughlin, B.A., Nelson, D., Silver, I.A., Erecinska, M., Chesselet, M.F., 1998. Methylmalonate toxicity in primary neuronal cultures. *Neuroscience* 86, 279–290.
- McLaughlin, B., Pal, S., Tran, M.P., Parsons, A.A., Barone, F.C., Erhardt, J.A., Aizenman, E., 2001. p38 activation is required upstream of potassium current enhancement and caspase cleavage in thiol oxidant-induced neuronal apoptosis. *J. Neurosci.* 21, 3303–3311.
- Rosenberg, P.A., 1991. Accumulation of extracellular glutamate and neuronal death in astrocyte-poor cortical cultures exposed to glutamine. *Glia* 4, 91–100.
- Rosenberg, P.A., Aizenman, E., 1989. Hundred-fold increase in neuronal vulnerability to glutamate toxicity in astrocyte-poor cultures of rat cerebral cortex. *Neurosci. Lett.* 103, 162–168.
- Skaper, S.D., Facci, L., Milani, D., Leon, A., Toffano, G., 1990. Culture and Use of Primary and Clonal Neural Cells, vol. 2. Academic Press, San Diego, pp. 17–33.
- Skaper, S.D., Facci, L., Kee, W.J., Strijbos, P., 2001. Potentiation by histamine of synaptically mediated excitotoxicity in cultured hippocampal neurons: a possible role for mast cells. *J. Neurochem.* 76, 47–55.
- Speliotis, E.K., Hartnett, K.A., Blitzblau, R.C., Aizenman, E., Rosenberg, P.A., 1994. Comparison of the potency of competitive NMDA antagonists against the neurotoxicity of glutamate and NMDA. *J. Neurochem.* 63, 879–885.
- Stanciu, M., Wang, Y., Kentor, R., Burke, N., Watkins, S., Kress, G., Reynolds, I., Klann, E., Angiolieri, M.R., Johnson, J.W., DeFranco, D.B., 2000. Persistent activation of ERK contributes to glutamate-induced oxidative toxicity in a neuronal cell line and primary cortical neuron cultures. *J. Biol. Chem.* 275, 12200–12206.
- Walton, K.M., DiRocco, R., Bartlett, B.A., Koury, E., Marcy, V.R., Jarvis, B., Schaefer, E.M., Bhat, R.V., 1998. Activation of p38MAPK in microglia after ischemia. *J. Neurochem.* 70, 1764–1767.